Sharbat Ahmad Shahi: A Potent Unani Formulation for Mood Disorders

Afshan Khan¹, Aisha Siddiqui^{2*}, M.A Jafri³, Mohd Asif⁴

^{1,2,3,4} Dept. of Ilmul Advia, School of Unani Medical Education and Research, Jamia Hamdard, New Delhi, India.

^{2*} Corresponding Author: Aisha Siddiqui, afshankhan2490@gmail.com

Available online at: www.isroset.org

Received: 10/Dec/2018, Accepted: 22/Dec/2018, Online: 31/Dec/2018

Abstract-Mood disorders such as depression are the most ubiquitous disease amongst psychiatric disorders. According to the estimations of the World Health Organisation, depression will be the second leading cause of disability in 2020. Psychiatric disorders are the foremost problems in the world since relic. In Unani system of medicine, there is no any direct description of depression but it is described in detail under the heading of “*Amraze nafsaniya*”. This system has a gem of valuable single and compound formulations for psychiatric disorders as there are several *mufrad* (Single) drugs used to treat psychiatric disorders such as Aftimoon (*Cuscuta reflexa*), Badranjboya (*Nepeta hindostana*), Nilofar (*Nymphaea alba*), Gule surkh (*Rosa damascena*), Bisfaij (*Polypodium vulgare*) etc. Sharbat Ahmad Shahi is one of the polyherbal Unani preparations and it has been conventionally used in *Malankholia* (Melancholia), *Insanity*, *Mania* and for *saudavi amraaz*. It has been reported for its antidepressant potential in recent years.

Key Words: Antidepressant, Mood disorders, Sharbat Ahmad Shahi, Unani medicine.

I. INTRODUCTION

Mood disorders such as depression are the most prevalent diseases amongst psychiatric disorders and a leading cause for disability worldwide. Depression, officially termed major depressive disorder (MDD) ranks among the most rampant diseases worldwide. Depression is diagnosed by the method, mentioned in the Diagnostic Manual of Mental Disorders (DSM-IV), which is characterized by the following symptoms: (1) depressed or irritable mood, (2) decreased interest or loss of pleasure, (3) weight gain or loss, (4) insomnia or hypersomnia, (5) psychomotor retardation or agitation, (6) fatigue or loss of energy, (7) feelings of worthlessness or inappropriate guilt, (8) diminished ability to think or concentrate, (9) recurrent thoughts of death and suicide. These symptoms must manifest daily for at least 2 weeks. The World Health Organization anticipated that depression will be the second leading cause of infirmity in 2020 [1]. Recent epidemiological studies specify that severe forms of depression affect 2-5% of the population worldwide, and up to 20% are influenced by milder forms of the disease [2]. Major depression is a serious disorder of massive sociological and clinical relevance. In conventional system of medicine, treatment outcome of depression is suboptimal. The use of currently available antidepressants is confined by their side effects, slow response, and inadequate treatment efficacy. Full remission is difficult to be achieved. Depression, as such is not mentioned in Unani classical texts as an individual disease entity rather it is mentioned as a symptom or group of symptoms of *Malankholia* (Melancholia) in which the mental functions of the

individual are deranged leading to constant grief, fear and dubious aggression. The patient find himself/herself deserted and occupied by loneliness and some patients experience delusion and hallucination also [3]. In Unani medicine, psychiatric disorders are dealt in detail under the heading of “*Amraze nafsaniya*” where it is described by various symptoms of psychic faculty and their distortion due to the involvement of vitiated humours. Unani System of medicine believes in holistic accession to the prevention and treatment of diseases. It covers physical, mental and spiritual dimensions of an individual's health. Several drugs have been reported to have beneficial effects in psychiatric (mental) disorders. There are several *mufrad* (Single) drugs used to treat psychiatric disorders such as Aftimoon (*Cuscuta reflexa*), Badranjboya (*Nepeta hindostana*), Nilofar (*Nymphaea alba*), Gule surkh (*Rosa damascena*), Bisfaij (*Polypodium vulgare*) etc and many *murakkab* (compound formulations) such as Khamira Gaozaban, Sharbat Aftimoon, Sharbat Ahmad Shahi and Itrifal Kishneezi etc have been reported to have beneficial effect in psychiatric (mental) disorders. Sharbat Ahmed Shahi (SAS) is one of them. It is a compound formulation usually prescribed for the patients of depression, insanity, melancholia, mania and for black bile disorders (*Saudawi amraz*) in Unani system of medicine for a long time due to its minimum side effects as compared to its counter synthetic antidepressants [4][5][6].

SHARBAT:

The word Sharbat is derived from Persian “Sharbat”, and sherbet is from Turkish “Serbet”, both of which in turn come

from Arabic “Sharba” a drink, from “Shariba” to drink. Sharbat is an important invention of a renowned Unani physician. It was invented by ‘Hakeem Fesaghoras’. It is a sweet viscous liquid or those medicinal preparations which are made either by preparing the decoction from the plant, animal and mineral origin drugs or by taking juice of the fruits from different plants and mixed with Sugar and boiled to the required consistency (*Qiwam*) i.e one tar [6][177][178].

METHOD OF PREPARATION OF SHARBAT:

Dry herbal drugs are soaked overnight in water, amounting to 8 or 10 times the weight of the drugs. Next morning, they

INGREDIENTS OF SHARBAT AHMAD SHAHI [6] [22]

are boiled till one third water is left, allowed to cool, rubbed with hands, and filtered through a piece of fine cloth. Then, two or three times of sugar are added, and the mixture is boiled on a low fire to obtain the desired consistency of Sharbat [177].

SHARBAT AHMAD SHAHI:

Sharbat Ahmed Shahi (SAS) is one of the polyherbal compound formulations. It has been traditionally used for the treatment of *Malikholia* (melancholia), Mania, psychoneurosis and palpitation. It is beneficial in cerebral insufficiency, also used in chronic constipation [4][6].

S.No	Unani Name	Botanical Name	Parts Used	Quantity
1	Aftimoon	<i>Cuscuta reflexa</i> Roxb.	Dried leaves	24g
2	Badranjboya	<i>Nepeta hindostana</i> Roth.	Leaves	225g
3	Berge Gaozaban	<i>Borago officinalis</i> Linn.	Leaves	200g
4	Bisfaij	<i>Polypodium vulgare</i> Linn.	Seeds	75g
5	Sana makki	<i>Cassia angustifolia</i> Vahl.	Leaves	9g
6	Berge Banafsha	<i>Viola odorata</i> Linn.	Leaves	50g
7	Gule Surkh	<i>Rosa damascena</i> Mill.	Flower	40g
8	Gule Nilofer	<i>Nymphaea alba</i> Linn.	Flower	75g
9	Halela siyah	<i>Terminalia chebula</i> Retz.	Fruit	75g
10	Tukhme Balangu	<i>Lallemantia royleana</i> Benth.	Seeds	75g
11	Qand safaid	Sugar		4kg

II. METHOD OF PREPARATION OF SHARBAT AHMAD SHAHI

SAS will be prepared as per the procedure, mentioned in (*Qarabadeen-e-Majeedi*) and (*Bayaz-e-Kabeer Part II*). First of all, the dry ingredients (mentioned above in the table) are soaked in 8 litres of water overnight and boiled next morning till only one-fourth volume of water remains. The decoction is now rubbed with the hands, cloth-filtered, and re-filtered through a cotton wool layer spread out on a sieve into another vessel. Next, the white sugar and citric acid are added to the decoction which is again heated. And any impurity that collects on the surface is removed. When the *Qiwam* becomes concentrated, the sodium benzoate dissolved in a little water is added, and immediately after boiling, the *Qiwam* is taken off the fire and sieved through a fine muslin cloth. Arq-e-gulab is now added to the Sharbat Ahmad Shahi, which is preserved in bottles.

Dosage & administration

25-35ml with 125ml of Arq Gaozaban or water [6]

1-2 Tola with water [4]

Pharmacological actions [4][6]

- Cephalic tonic (*Muqawwi Dimagh*)
- Cardio tonic (*Muqawwi Qalb*)
- Purgative of black bile (*Mushile Sauda*)

Therapeutic Uses [4][6]

- Melancholia (*Malikhuliya*)
- Schizophrenia (*Junoon*)
- Mania
- Mental disorders due to excess of sauda
- Black bile disorders (*Saudavi amraaz*)
- Brain disorders (*Amraze Dimaghi*)
- Palpitation (*Khafkan*)

Pharmacological Studies

- Anxiolytic activity[179]
- Antidepressant activity[179]
- Schizophrenia[14]

Brief description of Ingredients of Sharbat Ahmad Shahi:

Halela Siyah (*Terminalia chebula* Retz.)

- **PART USED:** Fruit [24]
- **DOSE:** 5-10g [23]
- **PHARMACOLOGICAL ACTIONS:**
 - Analgesic (*Musakkin*)[23]
 - Antibilious (*Dafe safra*)[25]
 - Astringent (*Qabiz*)[9]
 - Blood purifier (*Musaffi Khoon*)[26]
 - Brain tonic (*Muqawwi Dimagh*)[23]
 - Cardio-tonic (*Muqawwi Qalb*)[24]
 - Diuretic (*Mudire Baul*)[24]
 - Eye tonic (*Muqawwi Basar*)[23]
 - Intestinal tonic (*Muqawwi Amaa*)[9]
 - Laxative (*Mulaiyyan*)[26]
 - Purgative (*Mushil*)[25]
 - Stomachic (*Muqawwi Meda*)[26]
- **THERAPEUTIC USES:**
 - Amenorrhoea (*Ehtebase Tams*)[24]
 - Brain weakness (*Zofe Dimagh*)[23]
 - Bronchial asthma (*Zeequn Nafs*)[25]
 - Cardiac weakness (*Zofe Qalb*)[24]
 - Constipation (*Qabz*)[25]
 - Cough (*Suaal*)[25]
 - Dementia (*Zofe Hafiza*)[9]
 - Dysentery (*Zaheer*)[25]
 - Dyspepsia (*Tukhma*)[37]
 - Eye weakness (*Zofe Basar*)[23]
 - Facial paralysis (*Laqwa*)[9]
 - Fever (*Humma*)[26]
 - Flatulence (*Nafkhe Shikam*)[25]
 - Hepatomegaly (*Azme Kabid*)[26]
 - Hiccup (*Fuwaq*)[26]
 - Melancholia (*Malikholia*)[9]
 - Piles (*Bawaseer*)[9]
 - Rheumatism (*Hudar*) ([26]
 - Splenomegaly (*Azme Tihal*)[26]
 - Urinary diseases (*Amraze Baul*)[26]
 - Vomiting (*Qai*)[25]
 - Weakness of GIT (*Zofe Meda wa Ama*)[23]
 - Worms[26]
- **PHARMACOLOGICAL STUDIES**
 - Anti arthritic[79]
 - Anti caries activity[80]

- Anti HSV-2[71]
- Antiaging[75]
- Antibacterial [69]
- Anticonvulsant[73]
- Antidepressant[28]
- Anti-diabetic[35]
- Antifungal[76]
- Anti-inflammatory[78]
- Antimicrobial[72]
- Antioxidant[27]
- Antipyretic[77]
- Antiulcerogenic[70]
- Cardioprotective[32]
- Hepatoprotective[74]
- Immunomodulatory [31]
- Neuroprotective[33]

Bisfajej (*Polypodium vulgare* Linn.)

- **PART USED:** Root[24]
- **DOSE:** 5-10g[7]
- **PHARMACOLOGICAL ACTIONS**
 - Analgesic (*Musakkin*)[88]
 - Antiepileptic (*Mane Sara*)[7]
 - Anti-inflammatory (*Muhallil*)[8]
 - Antipyretic (*Dafe Humma*)[88]
 - Cardio tonic (*Muqawwi Qalb*)[7]
 - Digestive (*Hazim*)[88]
 - Diuretic (*Mudire Baul*)[88]
 - Hypothermic[88]
 - Laxative (*Mulaiyyan*)[88]
 - Purgative of black bile & phlegm (*Mushile Sauda wa Balgham*)[23]
- **THERAPEUTIC USES**
 - Asthma (*Damaa*)[8]
 - Colitis (*Qulanj*)[23]
 - Epilepsy (*Sara*) [8][7]
 - Eruptions[7]
 - Flatulence (*Nafkhe shikam*)[7]
 - Haemorrhoids (*Bawaseer*)[7]
 - Joint pain (*Mafasil ka dard*) [8]
 - Leprosy (*Juzam*)[7]
 - Melancholia (*Malikhuliya*)[8,7]
 - Rheumatic disorders (*Amraze hudar*) [7]
- **PHARMACOLOGICAL STUDIES**
 - Anti-arthritic[90]

- Anti-inflammatory[89]
- Antimicrobial[89]
- Antioxidant[89]

Aftimoon (*Cuscuta reflexa* Roxb.)

- **PART USED:** Whole plant [23]
- **DOSE:** 3-5g [23]
- **PHARMACOLOGICAL ACTIONS**
 - Anodyne ([36])
 - Anthelmintic (*Qatila Deedan*)[36]
 - Anti-inflammatory (*Muhallil*) [24]
 - Aphrodisiac (*Muqawwi bah*)[36]
 - Carminative (*Kasir riyah*)[36]
 - Diaphoretic[91]
 - Diuretic (*Mudire Baul*)[23][36]
 - Emmenagogue (*Mudire Haiz*)[25]
 - Expectorant (*Munaffis*)[36]
 - Purgative (*Mushil*)[91]
 - Sedative (*Munavim*)[36]
- **THERAPEUTIC USES**
 - Brain disorders (*Dimaghi Amraz*)[7]
 - Diseases of spleen (*Amraze Tihal*)[7]
 - Epilepsy (*Sara*)[92]
 - Facial paralysis (*Laqwa*)[7]
 - Flatulence (*Nafakhe Shikam*) [91]
 - Intestinal worms (*Deedane Ama*)[24]
 - Jaundice (*Yarqan*)[36]
 - Melancholia (*Malikhuliya*)[24]
 - Muscles & joint pain[36]
 - Numbness (*Khadar*)[7]
 - Palpitation (*Khafqan*)[7]
 - Paralysis (*Faalij*)[7]
 - Schizophrenia (*Junoon*)[24]
- **PHARMACOLOGICAL STUDIES:**
 - Antibacterial[95]
 - Anti-fertility activity[94]
 - Anticonvulsant[94]
 - Cytotoxic[93]
 - Anxiolytic[92]
 - Anti-inflammatory[93]

Gul-e-Surkh(*Rosa damacena* Mill.)

PART USED: Flowers [10][11]

DOSE: 5-7g [56]

- **PHARMACOLOGICAL ACTIONS:**

- Appetizer (*Mushtahi*) [57]
- Astringent (*Qabiz*) [58]
- Carminative (*Kasire Riyah*) [58]
- Refrigerant [58]
- Cardiotonic (*Muqawwi Qalb*) [58]
- Laxative (*Mulaiyyan*) ([59]
- Expectorant(*Munaffis*) [57]
- Antipyretic (*Dafe Humma*) [59]
- Aphrodisiac (*Muqawwi Bah*) [59]

- **THERAPEUTIC USES:**

- Sore throat (*Khushunate Halq*) [58]
- Tonsillitis (*Warne Lauzatain*) [58]
- Uterine hemorrhage [58]
- Urticaria (*Shara*) [58]
- Aphthous ulcer [58]
- Abdominal and chest pain [60]
- Menstrual problems (*Illate Haiz*) [60]
- Digestive problems [60]
- Headache (*Suda*) [59]
- Toothache[59]
- Stomatitis (*Qula*)[59]
- Wound healing [59]
- Cough (*Suaal*) [59]
- Dementia (*Zofe Hafiza*) [83]
- Reflux oesophagitis [86]

- **PHARMACOLOGICAL STUDIES:**

- Analgesic [61][62]
- Antiaging [63]
- Anticonvulsant [64]
- Antibacterial [65][66][67]
- Antidepressant [69]
- Antidiabetic [70][71][72]
- Anti-HIV [73]
- Antihypertensive [74]
- Anti-inflammatory [75]
- Antioxidant [76][77][78][79]
- Antispasmodic [80]
- Antitussive [81]

Badranjboya(*Nepeta hindostana* Roth.)

PARTS USED: Whole plant [88][89][90][91][92][93]

DOSE: 5-7g [94][95][96]

2 Tola [97][98]

- **PHARMACOLOGICAL ACTIONS:**

- Exhilarant (*Muffarah*) [99][100][101]
- Heart tonic (*Muqawwi Qalb*) [96][99][100][102]

- Concoctive (*Munzije sauda*) [94][95][96]
- Gout (*Dafeh Naqrus*) [101]
- Astringent (*Qabiz*) [95][103]
- Blood purifier (*Musafikhoon*) [95][97][98]
- Purgative (*Mushil*) [95][104]
- Arthritis (*Dafeh wajaul mafasil*) [101][105]
- Resolvent (*Muhallil*) [98][99][100]
- Stomachic (*Muqawwi Meda*) [95][100][106]
- Brain tonic (*Muqawwi Dimag*) [95][107][108]
- Deobstruent (*Muffateh Sudad*) [95][109][110]
- Demulscent (*Mulattif*) [96][99][111]
- Calorific (*Musakhin*) [96][99][110]
- Breast inflammation (*Dafeh waeme pistan*) [112]
- Carminative (*Kasire riyah*) [94]
- Antipyretic (*Dafeh Bukhar*) [94][113]
- Antidote (*Tiryaq sammom*) [94]

➤ THERAPEUTIC USES:

- Depression/Anxiety (*Parashani, Gumm*) [96][99][100][104][114]
- Insomnia (*Sehar*) [95][104][115]
- Cough (*Suaal*) [98][99][100]
- Bronchitis (*Warme shoeb*) [95][112][104]
- Hiccup (*Hichki*) [100][104]
- Fever (*Humma*) [94][95]
- Irritable Bowel Syndrome, Colitis (*Warme Qoloon*) [95][99]
- Dyspepsia (*Sue Hazam*) [100][109]
- Asthma (*Usre Tanafus*) [95][98][99]
- Menstrual problems (*Illate Haiz*) [98]
- Fever (*Humma*) [104][112]
- Hypertension (*Fisharrud dam*) [94][104]
- Migraine (*Shaqqeeqa*) [96]
- Influenza (*common cold*) [99][110][116]
- Shock (*Gashi*) [95]
- Muscular pain (*Wajaul Uzlaat*) [97][101][110]
- Vertigo (*Duwaar*) [99][109]
- Eczema (*Narfarsi*) [99]
- Alzheimer's disease [99][100]
- Sharpen Memory (*Muqawwi Hafiza*) [101][109][117]
- Antibacterial and Antifungal effects (*Dafe jaraseem wa Dafe fitrasraat*) [99]
- Leprosy (*Juzaam*) [95][97][112]

➤ PHARMACOLOGICAL STUDIES:

- Anti-depressant [104]
- Anti-oxidant [118]
- Anti-spasmodic [106]
- Analgesic and Sedation [119]

- Anti-tumor [120]
- Anti-microbial [121]
- Effect on cholinergic receptor [108]

Berge Gaozaban (*Borago officinalis* Linn.)

PART USED: Leaves [10]

DOSE: 5-7g (Berge Gaozaban) [10]
3-5g (Gule Gaozaban) [10]

➤ PHARMACOLOGICAL ACTIONS:

- Cardi tonic (*Muqawwi Qalb*) [122][123][124]
- Brain tonic (*Muqawwi Dimagh*) [122][123][124]
- Refrigerant [122][123][124]
- Demulcent (*Mulattif*) [122][123][124]
- Emollient [122][123][124]
- Laxative (*Mulaiyyan*) [122][123][124]
- Expectorant (*Munaffis*) [122][123][124]
- Lithotriptic (*Mufattite Hasat*) [122][123][124]
- Diuretic (*Mudire Baul*) [125][126][127]
- Febrifuge [125][126][127]
- Antispasmodic (*Dafe Tashannuj*) [125][126][127]
- Diaphoretic (*Moarriq*) [125][126][127]
- Lactagogue [125][126][127]
- Nervine [125][126][127]
- Sudorific [125][126][127]
- Depurative [125][126][127]
- Anti-inflammatory (*Muhallil*) [125][126][127]

➤ THERAPEUTIC USES:

- Cough (*Suaal*) [122][123][124]
- Coryza (*Zukam*) [122][123][124]
- Pneumonia (*Zatur Riyah*) [122][123][124]
- Pleurisy (*Zatul janab*) [122][123][124]
- Tuberculosis (*Diq*) [122][123][124]
- Asthma (*Zeequn Nafas*) [122][123][124]
- Meningitis (*Sarsam*) [122][123][124]
- Melancholia (*Malikhuliya*) [122][123][124]
- Functional palpitation of Heart [122][123][124]
- Jaundice (*Yarqaan*) [122][123][124]
- Stomatitis (*Qula*) [122][123][124]
- Kidney & Urinary bladder stones [122][123][124]
- Cancer (*Sartan*) [125]
- Corns [125]
- Sclerosis [125]
- Tumours (*Sala*) [125]
- Bronchitis (*Warme Shoeb*) [125]
- Fever (*Humma*) [125]
- Sore throat (*Khushunate Halq*) [125]
- Swellings & Urogenital ailments [125]

➤ **PHARMACOLOGICAL STUDIES:**

- Cardiovascular activity[13]
- Respiratory activity[13]
- Gastrointestinal activity[13]
- Antioxidant activity[180][181][182]
- Antibacterial activity[183]

Berg-e-Banafsha(*Viola odorata* Linn.)**PART USED:** Leaves and flowers [10][11]**DOSE:** Powder 3-6g [10]

Decoction 10-20ml [10]

➤ **PHARMACOLOGICAL ACTIONS:**

- Anti-inflammatory (*Muhallil*) [127][129]
- Diaphoretic (*Moarriq*) [126][127]
- Diuretic (*Mudire Baul*) [127][129]
- Emollient [126][127]
- Expectorant (*Munaffise Balgham*) [127]
- Laxative (*Mulaiyyan*) [126][129]
- Demulcent (*Mulattif*) [127]
- Emetic [127]
- Purgative (*Mushil*) [127]

➤ **THERAPEUTIC USES:**

- Cancer of Lungs, Breast or digestive tract [127]
- Whooping cough (*Shaheeqa*) [127]
- Headache (*Suda*) [127]
- Migraine (*Shaheeqa*) [127]
- Insomnia (*Seher*) [127]
- Bronchitis (*Warme Shoeb*) [127]
- Respiratory catarrh [127]
- Asthma (*Zeequn Nafas*) [127]
- Mouth & throat infections [127]
- Skin diseases [127]

➤ **PHARMACOLOGICAL STUDIES:**

- Antioxidant [128]
- Analgesic [129]
- Anti-inflammatory [130]
- Antipyretic [131]
- Sedative [132]
- Hypotensive & Lipid lowering effect [133]
- Antibacterial [134]
- Antihelmintic [135]
- Antifungal [136]
- Mosquito repellent activity [137]

Sana makki (*Cassia angustifolia* Vahl.)**PART USED:** Leaves and root [10][11]**DOSE:** 15-30mg [10]

500mg-2gm (Powder of leaf or pod) [10]

➤ **PHARMACOLOGICAL ACTIONS:**

- Laxative (*Mulaiyyan*) [138]
- Ophthalmic [138]
- Liver tonic (*Muqawwi Jigar*) [138]
- Cardiotonic (*Muqawwi Qalb*) [138]
- Expectorant (*Munaffise Balgham*) [138]
- Febrifuge [138]
- Antiasthma [139]
- Diuretic (*Mudire Baul*) [139]
- Also improve the visual activity [139]
- Purgative (*Mushil*) [140]

➤ **THERAPEUTIC USES:**

- Leprosy (*Juzam*) [141][142][143]
- Ringworm (*Quba*) [141][142][143]
- Flatulence (*Nafkhe Shikam*) [141][142][143]
- Colic (*Qulanj*) [141][142][143]
- Dyspepsia (*Sue Hazm*) [141][142][143]
- Constipation (*Qabz*) [141][142][143]
- Cough (*Suaal*) [141][142][143]
- Bronchitis (*Warme Shoeb*) [141][142][143]
- Cardiac disorders [141][142][143]
- Rheumatic disease (*Amraze Hudar*) [141][142][143]
- Leucoderma (*Bars*) [144][145]
- Eczema (*Narfarsi*) [144][145]

➤ **PHARMACOLOGICAL STUDIES:**

- Hypolipidemic [147]
- Antimutagenic [148]
- Hepatoprotective [150]
- Anti-inflammatory [150]
- Antishigellosis [149]
- Antibacterial [151]
- Antioxidant [155]
- Antiulcer [152]
- Antifungal [153, 154]
- Antitumour [146]
- Antifertility [146]

Gul-e-Nilofer (*Nymphaea alba* Linn.)**PART USED:** Rhizome and Flowers [11]**DOSE:** Flowers-30g [8]

➤ **PHARMACOLOGICAL ACTIONS:**

- Blood purifier (*Musafikhoon*) [156]
- Aphrodisiac (*Muqawwi Bah*) [156]
- Anodyne [156]
- Astringent (*Qabiz*) [156]
- Cardiotonic (*Muqawwi Qalb*) [156]
- Demulcent (*Mulattif*) [156]

➤ **THRAPEUTIC USES:**

- Insomnia (*Seher*) [156]
- Anxiety [156]
- Sore throat (*Khushunate Halq*) [156]
- Uterine cancer [156]
- Diaphoresis [157]
- Liver disorder [158]
- Urinary disorder [158]
- Menstrual problem (*Illate Haiz*) [158]
- Diabetes (*Ziabetes*) [158]
- Dyspepsia (*Sue Hazm*) [167]
- Heart problem [13][50][59][68]
- Cough (*Suaal*) [13][50][59][68]

➤ **PHARMACOLOGICAL STUDIES:**

- Antidiabetic [159]
- Hepatoprotective [160]
- Hyperproliferative response and renal carcinogenesis [161]
- Mutagenic activity [162]
- Anti proliferative [163]
- Tumor inhibitory activity [164]
- Cholinergic activity [164]
- Analgesic [164]
- Anti-inflammatory [164]
- Antiuro lithiatic [165]
- Uterotonic [166]
- Antimicrobial [164]

Tukhme Balangu (*Lallemantia royleana* Benth)

PART USED: Seeds [10][11]

DOSE: 5-7g [10]

➤ **PHARMACOLOGICAL ACTIONS:**

- Diuretic (*Mudire Baul*) [168][169]
- Tonic (*Stimulant*) [168][169]
- Aphrodisiac (*Muqawwi Bah*) [168][169]
- Antitussive [168][169]

➤ **THERAPEUTIC USES:**

- Hepatic disorder [168][169]
- Renal disorder [168][169]
- Nervous disorder [168][169]
- Psychotic disease [170]
- Rheumatism (*Hudar*) [171]
- Joint pain [171]
- Osteoarthritis [171]
- Fever (*Humma*) [172]
- Common cold [172]
- Expectorant (*Munaffis Balgham*) [172]
- Abscesses [17]

➤ **PHARMACOLOGICAL STUDIES:**

- Antibacterial activity [173]
- Hypocholesterolemic effect [174]
- Suspending agent [175]
- Thickening agent [175]
- Anesthetic activity [176]

III. CONCLUSION

This review article is endeavouring to bring out that Sharbat Ahmad Shahi is one of the best Unani formulations for psychiatric disorders. It has been reported to have beneficial effect in psychiatric (mental) disorders such as depression, insanity, anxiety etc. However more scientific studies and clinical trials are needed on this compound formulation to ensure its scientific validation for clinical use in patients.

REFERENCES

- [1]. Murray CJL and Lopez AD. The Global Burden of Disease: A comprehensive Assessment of Mortality & Disability from Disease, Injuries & Risk factors in 1990 & Projected to 2020. 1996 Cambridge, MA Harvard University Press Most comprehensive study of comparative disease burden even undertaken. Major depression estimated to be among the most burdensome disorders worldwide.
- [2]. R.C. Kessler and E. Bromet. The epidemiology of depression across cultures. Annu Rev Public Health. Author manuscript; available in PMC 2014 July 16.
- [3]. Anwar N, N.Z. Ahmed, T Shahida, K. Kabiruddin and H. Aslam. The Role of Mufarrehat (Exhilarants) in the Management of Depression: An Evidence Based Approach. J Psychiatry 20: 420.
- [4]. Kabiruddin H. Bayaze Kabir, Vol.2. Hyderabad; Hikmat Book Depot, 1938; pp. 80.

- [5]. Anonymous. National Formulary of Unani Medicine, Part I. New Delhi; Govt. Of India, Ministry of Health & Family Welfare, CCRUM, 2006; pp.85
- [6]. Anonymous. Qarabadeen-e-Majeedi. New Delhi; All India Unani Tibbi Conference: 1986, pp.197-198.
- [7]. Said HM. Hamdard Pharmacopoeia of Eastern Medicine. Delhi; Sri Satguru Publications, A division of Indian Books Centre, 1977; pp.290.
- [8]. Ghani MN, YNM. *Khazainul Advia* part III (Musawwer Edition). Idara Kitab-us-Shifa, New Delhi.pp. 985-988.
- [9]. Anonymous. Standardisation of single drugs of unani medicine, Part I,II, III, V. New Delhi; CCRUM, 1992; pp. 66-72, 241-146; 13-20,148-152, 282-288; 79-83; 16-24; 81-86; 214-217.
- [10]. Kabiruddin, M. H., 2000. *Makhzanul Mufridat Al Maroof Khawasul Advia*, Deoband and, Faisal Publication; pp.489,490,481,482,138,139,355,356,115,116.
- [11]. Hakeem, M. A., 2011. *Bustan Al Mufridat*. Musawwar Edition ed. New Delhi: Idara Kitabus-Shifa; pp.280,97,203,339,85.
- [12]. Gandhipuram PSK, Palanisamy A, Durairaj SK, Soremuthu PS. Antidiabetic activity of Fruits of Terminalia chebula on streptozoin induced diabetic rats. Journal of Health Science, 2006; 52 (3): 283–291
- [13]. Basu K. Indian Medicinal Plants, Vol. II: YNM, pp. 1741.
- [14]. Tahira P, Saida H, Nudrat A, Zubairi, Waseem A, Zehra B, Sumreen B. Effect of herbal combination on biochemical and behavioral responses in rats. Pak. J. Biochem. Mol. Biol. 2012; 45(1): 20-22
- [15]. Praveen S, Prakash T, Kotresha D, Asif MA, Uday RS, Bimlesh K, Jeevan D, Divakar G. Antitumorogenic activity of Terminalia chebula fruit in experimentally induced ulcer in rats. Pharmaceutical Biology, 2011 Mar; 49(3): 262-8
- [16]. Ajay K, Suja KP, Reshmi N, Aakanksha A, Nripendra NM, Satish KG. Anti-HSV-2 activity of Terminalia chebula Retz extract and its constituents, chebulagic and chebulinic acids. BMC Complementary and Alternative Medicine, 2017 17:110.
- [17]. Manoj K, Agarwal RC, Sanjay D, Rai VK, Benito J. Antimicrobial activity of aqueous extract of Terminalia chebula Retz on gram positive and gram negative microorganisms. International Journal of Current Pharmaceutical Research, 2009; 1(1)
- [18]. Jiban D, Uday RS, Bimlesh K, Nitesh SC. Anticonvulsant activity of ethanolic extract of fruits of Terminalia chebula on experimental animals. International Journal of Drug Development & Research, 2010;
- [19]. Mannan A, Khan RA, Asif M. Pharmacodynamic studies on Polypodium vulgare Linn. Indian J Exp Biol, 1989; 27(6): 556-60.
- [20]. Ambasta SP. The Useful plants of India. New Delhi; Publications & Information Directorate, CSIR, 1986;
- [21]. Anonymous. Unani Pharmacopoeia, Part I, Vol.I,II,III Vol. V. New Delhi; Department of AYUSH, Ministry of Health & Family Welfare, Government of India, 2007; 6, 18, 29, 32, 33, 106.
- [22]. Anonymous. Standardisation of single drugs of unani medicine, Part I,II, III, V. New Delhi; CCRUM, 1992; 1, 4, 11, 20, 24, 25, 79, 98-101,268,282-283,287.
- [23]. Khare CP. Indian Medicinal Plants. New Delhi; Springer: 2007, pp. 238, 365,652-653,654.
- [24]. Nadkarni KM. The Indian Materia Medica, vol. 1. Bombay; A.K Nadkarni Publishers: 1936, pp. 482,483, 842-843, 844-46.
- [25]. Hua-Yew C, Ta-Chen L, Kuo-Hua YU, Chien-Min Y, Chun-Ching L. Antioxidant and Free Radical Scavenging Activities of Terminalia chebula. Biol. Pharm. Bull, 2003; 26(9):1331-1335.
- [26]. Kabiruddin HM. *Makhzanul Mufridat Al Maroof Khawasul Advia*. Deoband; Faisal Publication: 2000. pp.55, 59,79-30, 74-75, 129-30, 145-146, 193-94,590-591.
- [27]. Nadkarni KM. Indian Plants & Drugs. New Delhi; Shrishti Book Distributors: YNM.pp. 151,393-394, 395-397.
- [28]. Chandra SR, Manohar VR, Rao SN. Antidepressant activity of aqueous extract of fruits of Terminalia chebula in rats. International Journal of Pharmacy and Pharmaceutical Sciences, 2012; 4(4): 449-451.
- [29]. Vaibhav A, Arun KW. Immunomodulatory Activity of Alcohol Extract of Terminalia chebula Retz Combretaceae. Tropical Journal of Pharmaceutical Research, 2011; 10(5): 567-575.
- [30]. Raman S, Srinivasan BP, Kiran D. Cardioprotective effect of Terminalia chebula in type II Diabetes induced cardiomyopathy in rat. World Journal of Pharmacy & Pharmaceutical Sciences, 2015; 4(10): 2151-2163.
- [31]. ChiaLin C, CheSan L. Phytochemical Composition, Antioxidant Activity and Neuroprotective Effect of Terminalia chebula Retz Extracts. Evidence-Based Complementary and Alternative Medicine, 2012; 1-7.
- [32]. Gandhipuram PSK, Palanisamy A, Durairaj SK, Soremuthu PS. Antidiabetic activity of Fruits of Terminalia chebula on streptozoin induced diabetic rats. Journal of Health Science, 2006; 52(3): 283-291.
- [33]. Praveen S, Prakash T, Kotresha D, Asif MA, Uday RS, Bimlesh K, Jeevan D, Divakar G. Antitumorogenic activity of Terminalia chebula fruit in experimentally induced ulcer In Rats. Pharmaceutical Biology, 2011; 49(3): 262-268.
- [34]. Ajay K, Suja KP, Reshmi N, Aakanksha A, Nripendra NM, Satish KG. Anti-HSV-2 Activity of Terminalia chebula Retz extract and its constituents, chebulagic and chebulinic acids. BMC Complementary and Alternative Medicine, 2017; 17(110): 1-11.
- [35]. Manoj K, Agarwal RC, Sanjay D, RAI VK, BENITO J. Antimicrobial activity of

- Aqueous Extract of Terminalia chebula Retz on gram positive and gram negative microorganisms. International Journal of Current Pharmaceutical Research, 2009; 1(1): 56-60.
- [36]. Jiban D, Uday RS, Bimlesh K, Nitesh SC. Anticonvulsant activity of ethanolic extract of Fruits of Terminalia chebula on experimental animals. International Journal of Drug Development & Research, 2010; 2(4): 764-768.
- [37]. Min-Kyung C, Hyeong-Geug K, Jong-Min H, Jin-Seok L, JongSuk L, SunHo C, Chang-GueSon. Hepatoprotective Effect of Terminalia chebula against t-BHP-Induced Acute Liver Injury in C57/BL6 Mice. Evidence-Based Complementary and Alternative Medicine, 2015; 1-11.
- [38]. Aranya M, Pensak J, Toshihiro A, Worapaka M, Jiradej M. In vitro anti-aging activities of Terminalia chebula gall extract. Pharmaceutical Biology, 2010; 48(4): 469-481.
- [39]. B. Rubini B, Shanthi G, Soundhari C, Rajarajan S. Antifungal activity of Terminalia chebula and Terminalia catappa on two dermatophytes. Journal of Medicinal and Aromatic Plants, 2013; 4(2): 15-19.
- [40]. Rajyalakshmi N, Nagesh RG, Ramanath B. Evaluation of local anaesthetic and antipyretic activity of Terminalia chebula Retz in experimental animal models. International Journal of Pharma and Bio Sciences, 2015; 6(1): 219-225.
- [41]. Saffath IJ, Zakia S, Ershad A, Mariam B, Mominul H. Evaluation of Analgesic and Anti-Inflammatory Activities on Ethanolic Extract of Terminalia chebula Fruits in Experimental Animal Models. American Journal of Plant Sciences, 2014; 63-69.
- [42]. Nichole M, Ramesh GC, Natalia V, Katie K, Jean M, John GT, Michelle LA, Terry CD, Sanyasi RK. Evaluation of Terminalia chebula Extract for Anti-Arthritic Efficacy And Safety in Osteoarthritic Dog. Veterinary Science & Technology, 2016; 7(1): 1-8.
- [43]. Carounanidy U, Satyanarayanan R, Velmurugan A. Use of an aqueous extract of Terminalia chebula as an anticaries agent: A clinical study. Indian J Dent Res, 2007; 18(4): 152-155.
- [44]. Veena S, Manu S. Operculina turpethum as a panoramic herbal medicine: A Review. International Journal of Pharmaceutical Sciences and Research, 2012; 3(1): 21-25.
- [45]. Ghani N. Khazainul Advia. Delhi; Idara Kitabus Shifa, YNM, pp. 187-89, 493, 526, 792, 793.
- [46]. Hakeem MA. Bustanul Mufradat. New Delhi; Idara Kitabus Shifa, 2002; 60, 67-68, 72, 85-86, 90-91, 92, 347, 348.
- [47]. Gaamoune S, Nouioua W, Djaout O. Antioxidant, antimicrobial and anti-inflammatory activities of flavonoids and tannins extracted from Polypodium vulgare L. Asian Journal of Biochemical and Pharmaceutical Research, 2015; 5(4): 114-122.
- [48]. Mannan A, Khan RA, Asif M. Pharmacodynamic studies on Polypodium vulgare Linn. Indian J Exp Biol, 1989; 27(6): 556-60.
- [49]. Ambasta SP. The Useful plants of India. New Delhi; Publications & Information Directorate, CSIR, 1986; 152.
- [50]. Basu K. Indian Medicinal Plants, Vol. II: YNM, pp. 1741.
- [51]. Sujith T, Sapna S, Velmurugan C, Ashok BS. Evaluation of anxiolytic activity of whole plant of Cuscuta reflexa. World Journal of Pharmacy and Pharmaceutical Sciences, 2015; 4(8): 1245-1253.
- [52]. Pavan BU, Suggala VS, Chandrashekhara DU. Preliminary screening of Cuscuta reflexa stems for Anti inflammatory and cytotoxic activity. Asian Afrin et al. European Journal of Biomedical and Pharmaceutical Sciences www.ejbps.com 206 Pacific Journal of Tropical Biomedicine, 2012; 1303-1307.
- [53]. Malaya G, Upal KM, Dilipkumar P, Shiladitya B, Sumit C. Studies on brain biogenic amines in methanolic extract of Cuscuta reflexa Roxb and Corchorus olitorius Linn seed treated mice. Acta Poloniae Pharmaceutica-Drug Research, 2003; 60(3): 207-210.
- [54]. Wiart C. Ethnopharmacology of Medicinal Plants. Asia and the Pacific Humana Press, 2006; 143.
- [55]. Ibn Sina. Al Qanoon fil Tib (Urdu Translation by Kantoori GH) Vols. 1, part 2, part 3, New Delhi; Idara Kitabus Shifa, 2007, pp. 39-40, 57, 61, 83-84, 212, 82-92.
- [56]. Lubhaya R. Goswami bayanul advia. Vol. I. Goswami Pharmacy, Qasim Jan Street, Delhi; 1982. p. 295-297.
- [57]. Chatterjee A, Pakrashi SC. The treatise on Indian medicinal plants. Vol. II. Publication & Information Directorate, CSIR, New Delhi; 1992. p. 15-16.
- [58]. Nadkarni AK. Indian Material Medica. Vol. I. Bombay Popular Prakashan, Mumbai; 1954. p. 1072-1073.
- [59]. Kirtikar KR, Basu BD. Indian Medicinal Plants. Vol. II. Connaught Place, Dehradun; 1991. p. 1072-1073.
- [60]. Boskabady MH, Shafei MN, Saberi Z, Amini S. Pharmacological effects of Rosa damascena. Iranian Journal of Basic Medical Sciences. 2011; 14(4): 295-307.
- [61]. Rakhshandeh H, Vahdati-Mashhadian N, Dolati K, Hosseini M. Antinociceptive effect of Rosa damascena in mice. Journal of Biological Sciences. 2008; 8(1): 176-180.
- [62]. Hajhashemi V, Ghannadi A, Hajiloo M. Analgesic and anti-inflammatory effects of Rosa damascena hydro alcoholic extract and its essential oil in animal models. Iranian Journal of Pharmaceutical Research. 2010; 9(2): 163-168.
- [63]. Jafari M, Zarban A, Pham S, Wang T. Rosa damascena decreased mortality in adult Drosophila. Journal of Medicinal Food. 2008; 11(1): 9-13.
- [64]. Kheirabadi M, Moghimi A, Rakhshandeh H, Rassouli MB. Evaluation of the

- anticonvulsant activities of *Rosa damascena* on the PTZ induced seizures in wistar rats. *Journal of Biological Sciences*.2008; 8:426-430.
- [65]. Kurhade BB, Vite MH, Nangude SL. Antibacterial activity of *Rosa damascena* Miller. *International Journal of Research in Pharmaceutical and Biomedical Sciences*. 2011; 2(3):1015-1020
- [66]. Ulusoy S, Boşgelmez-Tinaz G, Seçilmiş-Canbay H. Tocopherol, carotene, phenolic contents and antibacterial properties of rose essential oil, hydrosol and absolute. *Current Microbiology*. 2009; 59(5):554 -558.
- [67]. Halawani EM. Antimicrobial activity of *Rosa damascena* petals extracts and chemical composition by gas chromatography-mass spectrometry (GC/MS) analysis. *African Journal of Microbiology Research*. 2014; 8(24):2359-2367.
- [68]. K.R. Kritkar & B.D. Basu, *Indian Medicinal Plants*, International Book Distributors Dehradun 2005, vol 1, P.205-206.
- [69]. Dolati K, Rakhshandeh H, Shafei MN. Antidepressant-like effect of aqueous extract from *Rosa damascena* in mice. *Avicenna Journal of Phytomedicine*. 2011; 1(2):91-97.
- [70]. Ju JE, Joo YH, Chung N, Chung SY, Han SH, Lee YK. Anti-diabetic effects of red rose flowers in streptozotocin-induced diabetic mice. *Journal of Korean Society for Applied Biological Chemistry*.2014; 57(4):445-448.
- [71]. Gholamhoseinian A, Fallah H, Sharifi-far F. Inhibitory effect of methanol extract of *Rosa damascena* Mill. flowers on α -glucosidase activity and postprandial hyperglycemia in normal and diabetic rats. *International Journal of Phytotherapy and Phytomedicine*.2009;16(10):935-941.
- [72]. Gholamhoseinian A, Fallah H, sharifi-far F, Mirtajaddini M. The inhibitory effect of some Iranian plants extracts on the α glucosidase. *Iranian Journal of Basic Medical Sciences*. 2008; 11(1):1-9.
- [73]. Mahmood N, Piacente S, Pizza C, Burke A, Khan AI, Hay AJ. The anti-HIV activity And mechanisms of action of pure compounds isolated from *Rosa damascena*. *Biochemical and Biophysical Research Communication*.1996;229(1):73-79.
- [74]. Kwon EK, Lee DY, Lee H, Kim DO, Baek NI, Kim YE, Kim HY. Flavonoids from the buds of *Rosa damascena* inhibit the activity of 3-hydroxy-3-methylglutaryl-coenzyme A reductase and angiotensin I-converting enzyme. *Journal of Agricultural and Food Chemistry*. 2010; 58(2):882-886.
- [75]. Maleev A, Neshtev G, Stoianov S, Sheikov N. The ulcer protective and antiinflammatory effect of Bulgarian rose oil. *Eksperimental na Meditsinai Morfologiya*.1972; 11(2):55-60.
- [76]. Kumar N, Bhandari P, Singh B, Bari SS. Antioxidant activity and ultra-performance LC- electrospray ionization-quadrupole time-of-flight mass spectrometry for phenolics-based fingerprinting of rose species: *Rosa damascena*, *Rosa bourboniana* and *Rosa brunonii*. *Food and Chemical Toxicology*.2009; 47(2):361-367.
- [77]. Ozkan G, Sagdiç O, Baydar NG, Baydar H. Antioxidant and antibacterial activities of *Rosa damascena* flower extracts. *Food Science and Technology International*. 2004; 10(4):277-281.
- [78]. Yassa N, Masoomi F, Rohani Rankouhi SE, Hadjiakhoondi A. Chemical composition And antioxidant activity of the extract and essential oil of *Rosa damascena* from Iran, population of Guilan. *DARU Journal of Pharmaceutical Sciences*. 2009; 17(3):175-180.
- [79]. Shahriari S, Yasa N, Mohammadirad A, Khorasani R, Abdollahi M. In vivo antioxidant potential of *Rosa damascena* extract from Guilan, Iran comparable to α -tocopherol. *International Journal of Pharmacology*. 2007; 3(2):187-190.
- [80]. Bani S, Hasanpour S, Mousavi Z, Garehbaghi PM, Gojazadeh M. The effect of *Rosa damascena* extract on primary dysmenorrhea: a double-blind cross-over clinical trial. *Iranian Red Crescent Medical Journal*. 2014; 16(1):1-6.
- [81]. Shafei NM, Rakhshandeh H, Boskabady HM. Antitussive effect of *Rosa damascena* in Guinea pig. *Iranian Journal of Pharmaceutical Research*. 2003 ;(2): 231-234.
- [82]. Boskabady MH, Vatanprast A, Parsaee H, Ghasemzadeh M. Effect of aqueous-ethanolic extract from *Rosa damascena* on guinea pig isolated heart. *Iranian Journal of Basic Medical Sciences*. 2011; 14(2):116-121.
- [83]. Awale S, Tohda C, Tezuka Y, Miyazaki M, Kadota S. Protective effects of *Rosa damascena* and its active constituent on AB (25–35)-induced neuritic atrophy. *Evidence Based Complementary and Alternative Medicine*. 2011; (2011); Article ID 131042: 1-8.
- [84]. Rakshanda H, Hosseini M. Potentiation of phenobarbital hypnosis by *Rosa damascena* in mice. *Indian Journal of Experimental Biology*. 2006; 44(11):910-912.
- [85]. Arezoomandan R, Kazerani HR, Behnam-Rasooli M. The laxative and prokinetic effects of *Rosa damascena* Mill in rats. *Iranian Journal of Basic Medical Sciences*. 2011; 14(1): 9-16.
- [86]. Sengottuvelu S, Srinivasan D, Ramasamy S. The Effect of polyherbal formulation-PHF on experimentally induced reflux esophagitis in rats. *Journal of Pharmacy Research*. 2008; 1(1):11-15.
- [87]. Haleem, A., *Mufradatul Azizi* (Urdu), Matbua Sahitya Mandir Press Ltd., Lucknow 1948, p. 83, 86.
- [88]. Majoosi, A. *Kamilus Sanaah* (Urdu Translation by Ghulam Husain Kantoori), Matba Nami Munshi Naval Kishore 1889; V. 2, p. 313.
- [89]. Bolkent S, Yanardag R, Karabulut-Bulan O, et al. Protective role of *Melissa officinalis* L. extract on liver of hyperlipidemic rats: a morphological and biochemical study. *Journal of ethnopharmacology* 2005; 99:391-398
- [90]. Ibn Baitar. *Al Jami ul Mufradatul Adviawal Aghzia*. (Urdu translation CCRUM). New Delhi: Ministry of Health and Family Welfare, Govt. of India 1999; Vol. 3rd p.185.

- [91]. Ibn Kitabul Qanoon-fit-Tib (Translation), Matba Nawal Kishore, Lucknow, Edition 11037 H; p. 57.
- [92]. Kabeeruddin, H., (1924) *KulliyatAdvia*(Urdu), IdaraTameerTibb, Urdu Bazar, Lahore, p. 856-954.
- [93]. Samarqandi, N., *MualijateSharheAsbab*(Urdu translated by Kabeeruddin), Hikmat Book Depot, Hyderabad1916; V. 1, p. 156.
- [94]. Lubhaya, R. H., (NA) *Goswami Bayanul Advia*, Goswami Kutub Khana Gali Qasim Jaan, Delhi, V. 1, p. 122.
- [95]. Ibn Baitar *Aljamiul Mufradatal Adviaawal Aghzia* (Urdu translation), Central Council for Research in Unani Medicine, Ministry for Health and Family Welfare, New Delhi1985, V. 1, p. 177-179.
- [96]. Kabiruddin, H., () *BayazeKabir*(Urdu), Hikmat Book Depot, Hyderabad1935; pp. 313,218,189.
- [97]. Nadkarni K. M., *Indian Materia Medica*, Bombay Popular Prakashan, Bombay, India 1976; V.1, pp. 786.
- [98]. Hakim, M. A., (*BustanulMjufradat*(Urdu), KarkhanaJamitulAdvia, Lucknow, India1893; pp.83, 188.
- [99]. Haleem, A., *MufradateAzizi* (Urdu), MatbuaSahityaMandir Press Ltd., Lucknow 1948, pp.83, 86.
- [100]. Majoosi, A. *Kamilus Sanaah* (Urdu Translation by Ghulam Husain Kantoori), Matba Nami Munshi Naval Kishore1889; V. 2, pp. 313.
- [111]. Kabeeruddin, H., (1924) *KulliyatAdvia*(Urdu), IdaraTameerTibb, Urdu Bazar, Lahore, pp. 856-954.
- [112]. Samarqandi, N., *MualijateSharheAsbab*(Urdu translated by Kabeeruddin), Hikmat Book Depot, Hyderabad1916; V. 1, pp. 156
- [113]. Holmes JM. Thomas Jefferson treats himself: herbs, physicke, & nutrition in early America: Loft Press, Incorporated, 1997
- [114]. Perry N, Perry E. Aromatherapy in the management of psychiatric disorders: clinical And neuropharmacological perspectives. *CNS drugs* 2006; 20:257-280
- [115]. Bolkent S, Yanardag R, Karabulut-Bulan O, et al. Protective role of *Melissa officinalis* L.extract on liver of hyperlipidemic rats: a morphological and biochemical study. *Journal of ethnopharmacology* 2005; 99:391-398
- [116]. Kitano, Y., Usui, C., Takasuna, K., Hirohashi, M., Nomura, M., Increasing-current electroshock seizure test. A new method for assessment of anti-and proconvulsant activities of drugs in mice, *J PharmacoloToxicol methods*1996; V. 35, pp. 25-29.
- [117]. Gupta VK, Sharma SK. Plants as natural antioxidants. *Nat. Prod. Rad* 2006; 5:326-334.
- [118]. Mimica-Dukic N, Bozin B, Sokovic M, et al. Antimicrobial and antioxidant activities of *Melissa officinalis* L.(Lamiaceae) essential oil. *Journal of agricultural and food chemistry* 2004; 52:2485-2489
- [119]. Plat for a future, [<http://www.pfaf.org/User/plant.aspx?LatinName=Viola+odorata>]
- cited- 2017March 9.
- [120]. Dudai N, Weinstein Y, Krup M, et al. Citral is a new inducer of caspase-3 in tumor cell lines. *Planta Medica-Natural Products and Medicinal Plant Research* 2005; 71:484-487.
- [121]. Allahverdiyev A, Duran N, Ozguven M, et al. Antiviral activity of the volatile oils of *Melissa officinalis* L. against Herpes simplex virus type-2. *Phytomedicine* 2004; 11:657-661.
- [122]. Ghani N, Khazain ul advia, Vol 6.
- [123]. Hasan Masoda et al, 1982. *Jour Chem Soc Pak.*, 4(4): 281-283.
- [124]. Tewari NC et al, 1970. *Indian Jour Chem*, 8(7): 593-597.
- [125]. A handbook of Medicinal Herbs. James A Duke.
- [126]. Green Medicine. Leyel C.F, 93.
- [127]. Thoronton J, 1810. *New family of Herbal Medicine.*, 167.
- [128]. Ebrahimzadeh MA, Nabavi SM, Nabavi SF, Bahramian F, Bekhradnia AR. Antioxidant and free radical scavenging activity of *H. officinalis* L. var. *angustifolius*, *V. odorata*, *B. hyrcana* and *C. speciosum*. *Pak J Pharma Sci.* 2010; 23, 29-34.
- [129]. Barkatullah, Ibrar M, Ali N, Muhammad N, Meryam E. *In vitro* pharmacological study and preliminary phytochemical profile of *Viola canescens* Wall. *Ex Roxb. Afri J Pharm Pharmacol.* 2012; 6: 1142-46.
- [130]. Koochek MH, Pipelzadeh MH, Mardani H. The effectiveness of *Viola odorata* in the prevention and treatment of formalininduced lung damage in the rat. *J Herbs Spices Med Plants.*2003; 10: 95–103.
- [131]. Khattak SG, Gilani SN, Ikram M. Antipyretic studies on some indigenous Pakistani medicinal plants. *J Ethnopharmacol.*1985; 14: 45-51.
- [132]. Alireza M, Ali R. Evaluation of sedative and pre-anesthetic effects of *Viola odorata* Linn. extract compared with diazepam in rats. *Bull. Env. Pharmacol. Life Sci.* 2013; 2: 125-31.
- [133]. Siddiqi HS, Mehmood MH, Rehman NU, Gilani AH. Studies on the antihypertensive and antidyslipidemic activities of *Viola odorata* leaves extract. *Lipids Health Dis.* 2012;11: 6.
- [134]. Ramezani M, Zarrinkamar F, Bagheri M, Rajabnia, R. Study of environment temperature effect on the antibacterial activity of water extract of different organs of *Viola odorata* in the different stages of growth. *J Babol Univ Med Sci.* 2012; 14: 16–21.
- [135]. Colgrave ML, Kotze AC, Ireland DC, Wang CK, Craik DJ. The anthelmintic activity Of the cyclotides: Natural variants with enhanced activity. *Chem Bio Chem*, 2008, 9: 1939-45.
- [136]. Pawar V, Thaker V. *In vitro* efficacy of 75 essential oils against *Aspergillus niger*. *Mycoses* 2006; 49: 316-23.
- [137]. Amer A, Mehlhorn H. Repellency effect of forty-one essential oils against aedes, anopheles and culex mosquitoes. *Parasitol Res.* 2006; 99: 478–90.

- [138]. Hu S. Food Plants of China. Chinese University Press; 2005. p. 1–844.
- [139]. Chauhan NS. Medicinal and aromatic plants of Himachal Pradesh. Indus Pub. Co. New Delhi. 1999; 151-52)
- [140]. Manojlovic I, Bogdanovic-Dusanovic G, Gritsanapan W, Manojlovic N. Isolation and Identification of anthraquinones of *Caloplaca cerina* and *Cassia species*. Chemical Pap 2006; 60(6):466-68.
- [141]. Hooker, J.D., the Flora of British India, Vol.II, L.Reeve and Co., England, 1879, pp.26. Kirtikar, K.R. and Basu, B.D., Indian Medicinal Plants, Vol II, Periodical Experts D- 42, Vivek Vihar Delhi, 1975, pp.877.
- [142]. Jain, S.K., Medicinal Plants, National Book Trust, New Delhi., 1968, pp.37.
- [143]. Maitya TK, Mandal SC, Mukherjee PK. Saha K, Dass J, Saha BP, et al Evaluation of hepatoprotective potential of *Cassia species* leaf extract, Nat. Prod. Sci 1997; 3:122.
- [144]. Chauhan NS. Medicinal and aromatic plants of Himachal Pradesh. Indus Pub. Co. New Delhi. 1999;151-52.
- [145]. Anonymous. The Wealth of India A dictionary of Indian raw materials & Industrial products, Publications & Information Directorate, C.S.I.R., Vol.3, pp. 368-70.
- [146]. Anushia.C, Sampathkumar.P and Ramkumar.L. *Global Journal of Pharmacology*. 3 (3):127-130, 2009)
- [147]. Tiwari P, Kumar K, Panik R., Hepatoprotective effects of *Cassia species* whole plant. International Journal of Pharmacy & Technology 2011; 3(2):2798-2806.
- [148]. Patil UK, Saraf S, Dixit VK. Hypolipidemic activity of seeds of *Cassia species* Linn. J. of Ethnopharmacol 2004;90 (2-3):249-252.
- [149]. Choi JS, Lee HJ, Park KY, Ha JO, Kang SS. In vitro antimutagenic effects of anthraquinone aglycones and naphthopyrone glycosides from *Cassia species*. Planta Med 1997; 63 (1):11-14.
- [150]. Maitya TK, Mandal SC, Mukherjee PK. Saha K, Dass J, Saha BP, et al Evaluation of hepatoprotective potential of *Cassia species* leaf extract, Nat. Prod. Sci 1997; 3:122.
- [151]. Singh VK, Khan A M. Medicinal Plants and Folklores - A Strategy towards Conquest of Human Ailments. Vol. 9. Today & Tomorrow Printers & Publishers, 1990:67.
- [152]. Gill NS, Sharma A, Arora R, Bali M. Evaluation of *Cassia species* seeds for their antioxidant and antiulcer activity. J Med Sci 2011; 11 (2):96-101.
- [153]. Mukharjee PK, Saha K, Das J, Saha BP, Pal M. Antifungal activity of the leaf extract of *Cassia species* linn, Phytother. Res. 1996; 10:521-522.
- [154]. Acharya TK, Chatterjee IB, Isolation of chrysophanic acid –9-anthrone, a fungicidal compound from *Cassia species*, Sci.Cult.1974; 40:376.
- [155]. Chakrabarty K. Chawla H. Terpenoids and phenolics from *Cassia species* stem bark”, Indian J. Chem. 1983;22(B): 1165-1166.
- [156]. <http://www.pfaf.org/user/Plant.aspx?LatinName=Nymphaea+alba>
- [157]. Aspects of Asian medicine and its practice in the west by W.C Evans in Trease and Evans Pharmacognosy 15th edition W.B Saunders publications 2002, pp.no 477.
- [158]. Mohan Maruga Raja, M. K., Sethiya, Neeraj Kumar and Mishra S. H. (2010). A comprehensive review on *Nymphaea stellata*: A traditionally used bitter. J Adv Pharm Technol Res., 010 Jul-Sep; 1(3): 311–319.
- [159]. Rajagopal, S. and Sasikala, k. 2008. Antidiabetic activity of hydro-ethanolic extracts of *Nymphaea stellata* flowers in normal and alloxan induced diabetes rats. *Afri. J. Pharm. Pharmacol.* 2: 173-178.
- [160]. Bhandarkar, M.R. and Khan, A. 2004. Antihepatotoxic effect of *Nymphaea stellata* Willd. against carbon tetra chloride induced hepatic damage in albino rats. *J. Ethnopharmacol.* 91: 61-64.
- [161]. Khan, N. and Sultana, S. 2005. Chemomodulatory effect of *Ficus racemosa* extract against chemically induced renal carcinogenesis and oxidative damage response in Wistar rats. *Life Sci.* 29: 1194-1210.
- [162]. Sowemimo, A.A., Fakoya, F.A., Awopetu, I., Omobuwajo, O.R. and Adesanya, S.A. 2007. Toxicity and mutagenic activity of some selected Nigerian plants. *J. Ethnopharmacol.* 113: 427-432.
- [163]. Selvakumari, E., Shantha, S., Puroshothprabhu, T. And Sreenathkumar, C. 2012. Antiproliferative activity of Ethanolic flower extract from *Nymphaea pubescens* willd Youth Education and Research Trust (YERT) jairjp.com Selvakumari *et al.*, 2016 against human cervical and breast carcinoma *in vitro*. *Int. Res. J. Pharm.* 3: 124-125.
- [164]. Raja, M.M., Sethiya, N.K. and Mishra, S.H. 2010. A comprehensive review on *Nymphaea stellata*: A traditionally used bitter. *J. Adv. Pharmaceut. Technol. Res.* 3: 311-319.
- [165]. Bhaskar, V.H., Shelke, S. and Tushar, T. 2012. Effect of ethanolic extract of *Nymphaea alba* Linn on urolithiatic rats. *Int. J. Pharm. Pharmaceut. Sci.* 4: 572-574.
- [166]. Anindya, B., Moumita, S., Sudhanshu, S.R. and Sudam, C.S. 2014. Uterotonic properties of *Nymphaea alba* on Youth Education and Research Trust (YERT) jairjp.com Selvakumari *et al.*, 2016 isolated myometrium model. *Int. J. Pharm. Pharmaceut. Sci.* 6(6): 490-493.
- [167]. Sowemimo, A.A., Fakoya, F.A., Awopetu, I., Omobuwajo, O.R. and Adesanya, S.A. 2007. Toxicity and mutagenic activity of some selected Nigerian plants. *J. Ethnopharmacol.* 113: 427-432.
- [168]. Amin GR. 2005. Popular Medicinal Plants of Iran. Publications of Tehran University Of Medical Sciences, Tehran. 66-67.
- [169]. Ghannadi A, Zolfaghari B. 2003. Compositional analysis of the essential oil of

- Lallemantia royleana* (Benth. In Wall.) Benth. From Iran. Flavour Frag J, 18: 237-239.
- [170]. Safa O, Soltanipoor MA, Rastegar S, Kazemi M, Nourbakhsh Dehkordi Kh, Ghannadi A. 2013. An ethnobotanical survey on hormozgan province, Iran. Avicenna J Phytomed, 3: 64-81.
- [171]. Moghaddam TM, Razavi SMA, Emadzadeh B. Rheological interactions between *Lallemantia royleana* seed extract and selected food hydrocolloids. J. Sci. Food Agri, 2011; 91: 1083-1088.
- [172]. Naghibi F. Labiatae family in folk medicine in Iran. Iranian J. Pharm. Sci., 2005; (1): 200-210.
- [173]. Karcioğlu L, Tanis H, Comlekcioglu N, Diraz E, Kirecci E, Aygan A (2011). Antimicrobial Activity of *Salvia trichoclada* in Southern Turkey. Int. J. Agric. Biol. 13:134-136.
- [174]. Kanakavalli K, Thillaivanan S, Parthiban P, Vijayalakshmi G, Sudha M, Sutha J. 2014. Anti-hyperlipidemic herbs in siddha system of medicine. Int J Pharma Sci, 4: 541-545.
- [175]. Shanmygam S, Manavalan R, Venkappayya D. Natural polymer and their application. Natural product radiance, 2005; 4(6): 478-481.
- [176]. Rabi Atabaki and Majid Hassanpour-ezatti, Improvement of Lidocaine Local Anesthetic Action Using *Lallemantia royleana* Seed Mucilage as an Excipient. Iranian Journal of Pharmaceutical Research (2014), 13 (4): 1431-1436.
- [177]. Kabeeruddin M. Bayaz kabeer Part 3. New Delhi: Idara Kitabush Shifa pp 71-77, 2010.
- [178]. National formulary of Unani Medicine, Part –VI. Govt. of India, Ministry of Health & Family Welfare, CCRUM; 2011: 116.
- [179]. Muhammad A, Aisha A. Decreased brain serotonin turnover rate following administration of Sharbat-e-Ahmed Shah produces antidepressant effect in rats. Springer Science+Business Media, LLC 2017, Metab Brain Dis (2017) 32:1785–1790.
- [180]. B. Mhamdi, W. Aidi Wannes, J. Sriti, I. Jellali, R. Ksouri, B. Marzouk, “Effect of harvesting time on phenolic compounds and antiradical scavenging activity of *Borago officinalis* seed extracts”, Industrial Crops and Products, 2010, Vol. 31(1); 1-4.
- [181]. Mikel García-Iñiguez de Ciriano, Cecilia García-Herreros, Eduardo Larequi, Idoia Valencia, Diana Ansorena, Iciar Astiasarán, “Use of natural antioxidants from lyophilized water extracts of *Borago officinalis* in dry fermented sausages enriched in ω -3 PUFA”, Meat Science, 2009, Volume 83(2); 271-277.
- [182]. Mahinda Wettasinghe, Fereidoon Shahidi, “Antioxidant and free radical-scavenging properties of ethanolic extracts of defatted borage (*Borago officinalis* L.) seeds”, Food Chemistry, 1999, Vol. 67(4); 399-414.
- [183]. CEYLAN, E. and FUNG, D.Y.C. 2004. Antimicrobial activity of spices. J. Rapid Meth. Aut. Mic. 12, 1–55.

AUTHORS PROFILE

Afshan Khan presently pursuing M.D in Pharmacology at Jamia Hamdard University New Delhi, India. Mrs Aisha Siddiqui is currently working as Assistant Professor in Department of Ilmul Avia at Jamia Hamdard University New Delhi, India. Mr M. A Jafri is currently working as Professor in Department of Ilmul Advia at Jamia Hamdard University New Delhi, India and Mr Mohd Asif is currently working as Assistant Professor in Department of Ilmul Advia at Jamia Hamdard University New Delhi, India.